

L_1 COMPACTNESS OF BOUNDED BV SETS

ISIDORE FLEISCHER

ABSTRACT. Functions, uniformly bounded in BV norm in some bounded open set U in R^n , are compact in $L_1(U)$. This result is known when U has Lipschitz boundary [EG Th. 4 p. 176], [G 1.19 Th. p. 17], [Z 5.34 Cor. p. 227]; the proof for general U here, after identifying the operator theoretic definition of bounded BV norm with that of the Tonelli variation, appeals to the standard compactness criterion in L_1 [DS 21 TH. p. 301] [Y, p. 275] (For completeness, these two auxiliary results are also presented).

PROOF

For the purpose of establishing the compactness criterion, it will be necessary to bound integrals of the form $\int_0^1 V(x+h) - V(x) dx$ for monotone non-decreasing V on $[0, 1]$, constant on $[1, \infty)$, for $h > 0$. Since $\int_0^1 V(x+h) dx = \int_h^{1+h} V(x) dx$, the difference integrates to $\int_1^{1+h} V(x) dx - \int_0^h V(x) dx$, bounded by $2h[V(1) - V(0)]$. It thus appears that the integral of the differences goes to 0 with h , uniformly in the rise $V(1) - V(0)$ of V . If V is the variation of a BV function F , then the same bound holds for $\int |F(x+h) - F(x)| dx$. If the V are functions of other variables, one can also obtain this uniformity for the integrals extended over bounded (in measure) subsets of these.

THE COMPACTNESS CRITERION

A bounded subset $\mathcal{K}$ of $L_1(U)$ (U open bounded), satisfying $\lim_{t \rightarrow 0} \int |f(s+t) - f(s)| ds = 0$ uniformly for $f \in \mathcal{K}$ is precompact.

The hypothesis entails (by $s \rightarrow s+t_0$) the formally stronger $\int |f(s+t) - f(s+t_0)| ds \rightarrow 0$ for every t_0 and uniformly in t_0 as well as in $\mathcal{K}$.

$(M_r f)s := \int_{B_r} f(s+t) dt$ ($f_E := (1/|E|) \int$, average of integrand over measurable E ; $B_r :=$ centered ball of radius r) is uniformly equicontinuous in $\mathcal{K}$ and $r \geq r_0 > 0$ and has the same L_1 norm as f – hence, this set of $M_r f$ is precompact (in C , by Ascoli, hence) in L_1 .

$\|M_r f - f\| = \int |\int_{B_r} f(s+t) - f(s) dt| ds \leq \int_{B_r} \int |f(s+t) - f(s)| ds dt \leq \sup_{t \in B_r} \int |f(s+t) - f(s)| ds \rightarrow 0$ as $r \downarrow 0$, uniformly in $\mathcal{K}$. It thus suffices to verify that $\{M_r f : r > 0, f \in \mathcal{K}\}$ is precompact and for this, that every sequence with $r \rightarrow 0$ has a subsequence every two terms of which are $< 1/n$ apart. For any such sequence, find an r_0 below which $M_r f$ is within $1/2n$ of

f uniformly in $\mathcal{K}$ and use the precompactness of the $M_{r_0}f$ to extract such a subsequence. By diagonalizing, this will produce a Cauchy subsequence.

BV VARIETIES

The (more precisely, “A”) *variation* of an n - argument real-valued locally integrable function F in an open bounded set U is defined to be the sup of $\int_U F \nabla \cdot \varphi dx$ over the C' φ to $[-1 + 1]$ with compact support in U .

The integral is a sum of n terms $\int_U F \partial \varphi_i / \partial x_i dx_i d\tilde{x}_i$ where $d\tilde{x}_i$ is the element of $(n-1)$ -volume in the co-ordinate hyperplane orthogonal to the x_i -axis and φ_i is the i^{th} component of φ . Since φ could have just this component different from 0, the sup of the sum dominates the sup of each term; and the sum of these non-negative sups dominates the sup of their sum. Hence bounded variation comes to finiteness of the n sups of $\int_U F \partial \varphi_i / \partial x_i dx_i d\tilde{x}_i = \int_U -\varphi_i d_i F d\tilde{x}_i$ where $d_i F$ is the distribution in x_i obtained from F by fixing the other co-ordinates; the minus sign can be dropped since the φ 's are closed for negation. The value of the integral is unchanged by making F right continuous in x_i , so assume it so. The sup is dominated by the “Tonelli Variation” $\int V_x(F) d\tilde{x}$, the integrand being the classical variation of right continuous F with all the variables other than x fixed.¹ This quantity is actually attained by the sup as we now show.

It would suffice to attain an arbitrary approximating sum whose terms are measures of finitely many disjoint $\tilde{x}$ -measurable subsets with values of the integrand at points in the sets as coefficients: this integrand is itself a sup of approximating sums: $\Sigma |F(t_{i+1}) - F(t_i)|$ with t_i points of continuity of F .

It is easy to find a sequence of $[-1, +1]$ -valued C' functions on a real interval $[s, t]$ vanishing, along with their derivatives, at s and t which increase pointwise to $1_{(s, t)}$; the integral of this sequence against $BV dF$ converges to $F(t^-) - F(s^+)$. Glueing together translates of this sequence or its negative yields a C' sequence to $[-1, +1]$ whose integral against dF converges to $\Sigma |F(t_{i+1}) - F(t_i)|$ provided the partition points t_i are restricted to points of continuity of F .

One can also find a sequence of $[-1, +1]$ -valued C' functions on $\tilde{X}$ which converges a.e. to the characteristic function of a given measurable set: represent the set and its complement a.e. as a union of a sequence of closed sets; for like-indexed pairs find (by Urysohn) a continuous function 1 on the contained closed set and 0 on the one disjoint; and approximate the resulting sequence (by Weierstrass-Stone) by a C' sequence with the same pointwise limit. The $\tilde{x}$ -integral of these functions converges to the measure of the set (which the function characterizes) and so their pairwise product with the real (x -) variable sequence previously constructed yields a C' sequence on

¹The integrand is positive measurable as limit of a sequence of such; a priori the integral could be infinite.

$\tilde{X} \times X$ to $[-1, +1]$ whose $\int \varphi dF dx$ converges to $\Sigma |F(t_{i+l}) - F(t_i)|$ multiplied by the measure of the $\tilde{x}$ -subset, which is one term of the sought for approximating sum; adding these finitely many sequences pointwise yields a sequence converging to the full sum.

P.S. The same result is valid with the usual (Vitali) notion of BV : Over a bounded domain, a uniformly BV , L_1 -bounded set of functions is uniformly bounded (for if the values at some point were unbounded then by BV uniformity so would be their lower bounds uniformly over the domain; and then again by uniform BV , the pointwise bounded set would be uniformly bounded over the domain). By Helly's Selection Theorem, some subsequence converges pointwise, whence by the uniform boundedness, in L_1 .

REFERENCES

- [DS] N. Dunford and J. Schwartz, *Linear Operators*, Pt 1. Interscience.
- [EG] L. C. Evans and R. F., Gariepy, *Measure theory and fine properties of functions*, CRC Press, 1992.
- [G] E. Giusti, *Minimal Surfaces and Functions of Bounded Variation*, Birkhäuser, 1984.
- [Y] K. Yosida, *Functional Analysis*, Springer Verlag, 1968.
- [Z] W. P. Ziemer, *Weakly differentiable functions*, Springer-Verlag 1989.

CENTRE DE RECHERCHES MATHÉMATIQUES, C.P. 6128, SUCCURSALE CENTRE-VILLE,
MONTRÉAL, QC, H3C 3J7, CANADA